

POST-OPERATIVE INSTRUCTIONS

Sometimes the after-effects of oral surgery are quite minimal, so not all of these instructions may apply. Common sense will often dictate what you should do. However, when in doubt, follow these guidelines or call our office for clarification. Our number is **845-340-1962**. Please visit our website www.midhudsonoralsurgeon.com

DAY OF SURGERY

FIRST HOUR: Bite down gently but firmly on the gauze packs that have been placed over the surgical areas, making sure they remain in place. Do not change them for the first two hours unless the bleeding is not being controlled. If active bleeding persists after two hours, place enough new gauze to obtain pressure over the surgical site for another 45-60 minutes. The gauze may be changed as necessary (typically every 30 to 45 minutes). It is best to moisten the gauze with tap water and loosely fluff for more comfortable positioning.

EXERCISE CARE: Do not disturb the surgical area today. **DO NOT rinse vigorously** or probe the area with any objects or your fingers. You may brush your teeth gently. **PLEASE DO NOT SMOKE** for at least 48 hours, since it is very detrimental to healing, and may cause a dry socket.

OOZING: Intermittent bleeding or oozing is normal; it may be controlled by placing fresh gauze over the surgical areas and biting down firmly for 30-60 minutes at a time.

PERSISTANT BLEEDING: Bleeding should never be severe. If it is, it usually means that the packs are being clenched between your teeth only, rather than exerting pressure on the surgical areas. Try repositioning the gauze packs. If bleeding persists or becomes heavy, you may substitute the gauze with a tea bag (soaked in hot water, squeezed damp-dry and wrapped in a moist gauze) for 20-30 minutes. ***If bleeding remains uncontrolled, please call our office.***

SWELLING: Swelling is often associated with oral surgery. Using a cold pack can minimize it or ice bag wrapped in a towel and applied firmly to the face or cheek adjacent to the surgical area. The ice pack should be applied twenty minutes on and twenty minutes off during the first 24 hours after surgery. If you have been prescribed medicine for the control of swelling, be sure to take it as directed.

PAIN: Unfortunately most oral surgery is accompanied by some degree of discomfort. You will usually have a prescription for pain medication. If you take the first pill before the local anesthetic has worn off, you will be able to manage any discomfort better. Effects of pain medications vary widely among individuals. If you do not achieve adequate relief, you may supplement each pill with an analgesic such as acetaminophen. Some people may require two pain medication pills at one time during early stages following surgery (**but that may add to the risk of upset stomach**). **Remember that the most severe discomfort is usually within the first six hours after the anesthetic wears off, after that your need for medication should lessen.**

NAUSEA Nausea is not an uncommon event after surgery, and it is sometimes caused by stronger pain medications. Nausea may be reduced by preceding each pill with a small amount of soft food, then taking the pill with a large volume of water. Try to keep taking clear fluids and minimize the pain medication, but call us if you do not feel better or if repeated vomiting is a problem. Cola drinks that have less carbonation may help with nausea.

DIET Eat any nourishing food that can be taken with comfort. Temperature of the food doesn't matter, but avoid extremely hot foods. It is sometimes advisable (but not required) to confine the first day's intake to bland liquids or pureed foods (creamed soups, puddings, yogurt, milk shakes, etc). Avoid foods like nuts, sunflower seeds, popcorn, etc. that may get lodged in the socket areas. Over the next several days you can progress to solid foods at your own pace. It is important not to skip meals! If you take nourishment regularly, you will feel better, gain strength, have less discomfort and heal faster. If you are a diabetic, maintain your normal eating habits as much as possible and follow the instructions given to you by our office or your physician regarding your insulin schedule.

SHARP EDGES If you feel sharp edges in the surgical areas with your tongue it is probably the bony wall which originally supported the teeth. Occasionally small slivers of bone may work themselves out during the first week or two after surgery. They are not pieces of tooth and if necessary, we will remove them. Please call the office if you are concerned.

INSTRUCTIONS FOR THE SECOND AND THIRD DAYS

MOUTH RINSES

Keeping your mouth clean after surgery is essential. Use one-quarter teaspoon of salt dissolved in an 8 ounce glass of warm water and gently rinse with portions of the solution, taking five minutes to use the entire glassful. Repeat as often as you like, but at least two or three times daily for the next five days.

BRUSHING

Begin your normal oral hygiene routine as soon as possible after surgery. Soreness and swelling may not permit vigorous brushing of all areas, but please make every effort to clean your teeth within the bounds of comfort.

HOT APPLICATIONS

Apply warm compresses to the skin overlying areas of swelling (hot water bottle, moist hot towel, and heating pad) for 20 minutes on, 20 minutes off to help soothe those tender areas. This will also help decrease swelling and stiffness.

SYRINGE

If you were given an irrigation syringe at your first office check-up visit, start using it the third day after surgery to keep the sockets clean. Fill it with warm water and irrigate any open sockets gently, especially after eating.

DRY SOCKETS

Normal healing after tooth extraction should be as follows: The first day of surgery is usually the most uncomfortable and there is some degree of swelling and stiffness. The second day you will usually be far more comfortable and, although still swollen, you can usually begin a more substantial diet. From the third day on **GRADUAL, STEADY IMPROVEMENT** should mark the remainder of your post-operative course. If a **DRY SOCKET** occurs (loss of blood clot from socket, usually on the 3rd to 5th day), there is a noticeable, distinct, persistent throbbing pain in the jaw, often radiating toward the ear and forward along the jaw to cause other teeth to ache. If you do not see steady improvement during the first few days after surgery, don't suffer needlessly. Call the office and report symptoms so you can be seen as soon as possible.

It is our desire that your recovery be as smooth and pleasant as possible. Following these instructions will

assist you, but if you have questions about your progress, please call the office where you had your surgery. Please try to call during office hours, however a 24-hour answering service is available for after hours contact with a doctor. Calling during office hours will allow for a faster response.